

Near vision from an eye drop? Yes.

USES: Qlosi (pilocarpine ophthalmic solution) 0.4% is a prescription eye drop used to treat age-related blurry near vision (presbyopia) in adults.

Blurry vision when trying to see up close can be frustrating.

Losing the ability to see up close and focus on small details (known as presbyopia) is a reality we all will most likely face. With this condition, you may need to rely on your readers or contacts to focus on things like reading on your phone or a menu.





Qlosi eye drops may help you achieve clear near vision, when you need it.

Imagine going to dinner and reading the menu without your readers. Moments like this are made for Qlosi—an FDA-approved eye drop that improves your eye's natural ability to focus on nearby objects and small details.

IMPORTANT SAFETY INFORMATION

 \cdot Do not use Qlosi if you are allergic to any of the ingredients

Qlosi[™] is comfortable and easy to use throughout the day.



Once or twice daily dosing It's up to you—after using one drop in each eye, you can use a second drop after 2 to 3 hours to extend your near vision (up to 8 hours).




Easy-to-use vials You can take Qlosi with you on the go.



Works with other eyedrops Wait 5 minutes after using Qlosi before using the other eye drops.



Okay if you wear contacts Simply remove contacts, use Qlosi, and wait 10 minutes before putting them back in.



<text>

No insurance approval or extra steps are needed. Qlosi can be purchased using personal funds or savings debit or credit. You can also pay using the funds in your Flexible Spending Account (FSA) or Health Savings Account (HSA).



Ask your eye doctor about Qlosi.



A low likelihood of side effects

Qlosi was studied in five US clinical trials with people who have presbyopia. The most common side effects (5% to 8%) were eye discomfort after first using the drops and headache. Most people did not experience side effects at all, and the side effects were mild, brief, and went away after a short time.

IMPORTANT SAFETY INFORMATION

- $\cdot\,$ Do not use Qlosi if you are allergic to any of the ingredients
- Use caution when driving at night or performing hazardous activities in poor lighting. Qlosi may cause temporary dim or dark vision.
- Temporary problems when changing focus between near objects and distant objects may occur. Do not drive or operate machinery if your vision is not clear
- Seek immediate medical care if you experience sudden onset of flashes of light, floaters or vision loss.
- If you wear contact lenses, they should be removed before using Qlosi. Wait 10 minutes after dosing before reinserting your contact lenses.
- Do not touch the tip of the vial to the eye or any other surface to prevent eye injury or contamination
- If you use more than one eye medication, use each medication at least 5 minutes apart
- The most common side effects reported with Qlosi in clinical trials are eye pain when administering Qlosi in your eye and headache. These are not all of the possible side effects with Qlosi.

To report suspected side effects, contact Orasis Pharmaceuticals at 1-866-ORASIS-1 or FDA at www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see full Prescribing Information located in the Qlosi package or visit glosi.com

It's possible to see up close with Qlosi™



Scan the QR code below or visit **Qlosi.com** to sign up for emails and learn more.



©2025 Orasis Pharmaceuticals Inc. All Rights Reserved. QLO.00119 03/2025